

***Centromadia parryi* (Greene) Greene**  
**ssp. *parryi***

PAPPOSE TARPLANT  
ASTERACEAE

Synonym: *Hemizonia parryi* ssp. *parryi*

**Status:**

CNPS Rare Plant Rank 1B.2  
Federal: unlisted  
State: unlisted  
State Rank: S1

**Habitat:** Often alkaline; chaparral, coastal prairie, meadows and seeps, marshes and swamps (coastal salt), valley and foothill grassland (vernally mesic). 0 - 420 meters (0 - 1,380 feet)



Toni Corelli

**Counties:**

Butte, Colusa,  
Glenn, Lake,  
Napa, **San**  
**Mateo**, Solano,  
Sonoma

**Life Form:**

Dicot  
annual herb

**Bloom period:**

May - November



[CNPS Rare and Endangered Plant Inventory Listing](#)

[Jepson eFlora Taxon page](#)

[Jepson eFlora key](#)

[Calflora Taxon Report](#)

[CalPhotos Database Results](#)

[Consortium of California Herbaria Records](#)



Toni Corelli



Aaron Schusteff

***Centromadia parryi* (Greene) Greene ssp. *parryi***



*Avis Boutell*

*Centromadia parryi* ssp. *parryi* (PAPPOSE TARPLANT)

Peduncle bracts coarsely glandular, glands sessile or short-stalked  
compare with CONGDON'S TARPLANT:



*Toni Corelli*

*Centromadia parryi* ssp. *congdonii* (CONGDON'S TARPLANT)

Peduncle bracts seldom glandular